

## Referenties – Tessa Scheffers-van Schayck, Karin Monshouwer & Marlous Tuithof

- [1] Oostingh, E.C., Hall, J., Koster, M.P.H., Grace, B., Jauniaux, E., & Steegers-Theunissen, R.P.M. (2018). The impact of maternal lifestyle factors on periconception outcomes: A systematic review of observational studies. *Reprod Biomed Online*, 38(1): 77-94. doi: 10.1016/j.rbmo.2018.09.015
- [2] Hofhuis, W., De Jongste, J.C., & Merkus, M. (2003). Adverse health effects of prenatal and postnatal tobacco smoke exposure on children. *Arch Dis Child.*, 88: 1086-90.
- [3] Scheffers-van Schayck, T., Thissen, V., de Josselin de Jong, S., Van der Houwen, C., Smith, S., & Steegers-Theunissen, R. (2021). Factsheet Rookvrij zwanger worden: Literatuuroverzicht van de risico's van (mee)roken in de periconceptie periode. Utrecht: Trimbos-instituut.
- [4] U.S. Department of Health and Human Services (2014). The health consequences of smoking – 50 years of progress: A report of the surgeon general. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.
- [5] Sundermann, A.C., Zhao, S., Young, C.L., Lam, L.A., & Jones, S.H. (2019). Alcohol use in pregnancy and miscarriage: A systematic review and meta-analysis. *Alcohol Clin Exp Res.*, 43(8): 1606-16. doi: 10.1111/acer.14124
- [6] Mamluk, L., Edwards, H.B., Savović, J., Leach, V., Jones, T., Moore, T.H.M., et al. (2017). Low alcohol consumption and pregnancy and childhood outcomes: Time to change guidelines indicating apparently “safe” levels of alcohol during pregnancy? A systematic review and meta-analyses. *BMJ Open.*, 7(7). doi: 10.1136/bmjopen-2016-015410
- [7] Caputo, C., Wood, E., & Jabbour, L. (2016). Impact of fetal alcohol exposure on body systems: A systematic review. *Birth Defects Res C Embryo Today.*, 108(2): 174-80. doi: 10.1002/bdrc.21129
- [8] U.S. Department of Health and Human Services. (2006). The health consequences of involuntary exposure to tobacco smoke: A report of the surgeon general. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.
- [9] Anderson, T.M., Lavista Ferres, J.M., You Ren, S., Moon, R.Y., Goldstein, R.D., Ramirez, J.M., et al. (2019). Maternal smoking before and during pregnancy and the risk of sudden unexpected infant death. *Pediatrics*, 143(4). doi: 10.1542/peds.2018-3325
- [10] Matt, G.E., Quintana, P.J.E., Destailats, H., Gundel, L.A., Sleiman, M., Singer, B.C., et al. (2011). Thirdhand tobacco smoke: Emerging evidence and arguments for a multidisciplinary research agenda. *Environ Health Perspect*, 119(9): 1218-26. doi: 10.1289/ehp.1103500
- [11] Giglia, R., & Binns, C. (2006). Alcohol and lactation: A systematic review. *Nutr Diet.*, 63(2): 103-16. doi: 10.1111/j.1747-0080.2006.00056.x
- [12] Gibson, L., & Porter, M. (2018). Drinking or smoking while breastfeeding and later cognition in children. *Pediatrics*, 142(2). doi: 10.1542/peds.2017-4266
- [13] Smedberg, J., Lupattelli, A., Mardby, A.-C., & Nordeng, H. (2014). Characteristics of women who continue smoking during pregnancy: A cross-sectional study of pregnant women and new mothers in 15 European countries. *BMC Pregnancy Childbirth*, 14(1): 213. <https://doi.org/10.1186/1471-2393-14-213>
- [14] Nur, N. (2017). Association of risk factors with smoking during pregnancy among women of childbearing age: An epidemiological field study in Turkey. *Sao Paulo Med J.*, 135(2): 100-6. doi: 10.1590/1516-3180.2016.021921102016
- [15] Harkonen, J., Lindberg, M., Karlsson, L., Karlsson, H., & Scheinin, N.M. (2018). Education is the strongest socio-economic predictor of smoking in pregnancy. *Addiction*, 113(6): 1117-1126. doi: 10.1111/add.14158
- [16] Mohsin, M., & Bauman, A.E. (2005). Socio-demographic factors associated with smoking and smoking cessation among 426,344 pregnant women in New South Wales, Australia. *BMC Public Health*, 5: 1-9. doi: 10.1186/1471-2458-5-138

- [17] Kia, F., Tosun, N., Carlson, S., & Allen, S. (2018). Examining characteristics associated with quitting smoking during pregnancy and relapse postpartum, 78: 114-9. doi: 10.1016/j.addbeh.2017.11.011
- [18] Lu, Y., Tong, S., & Oldenburg, B. (2001). Determinants of smoking and cessation during and after pregnancy. *Health Promot Int.*, 16(4): 355-65. doi: 10.1093/heapro/16.4.355
- [19] Scheffers-van Schayck, T., Tuithof, M., Otten, R., Engels, R., & Kleinjan, M. (2019). Smoking behavior of women before, during, and after pregnancy: Indicators of smoking, quitting, and relapse. *Eur Addict Res.*, 25: 132-44. doi: 10.1159/000498988
- [20] Shmulewitz, D., & Hasin, D.S. (2019). Risk factors for alcohol use among pregnant women, ages 15-44, in the United States, 2002 to 2017. *Prev Med.*, 124:75. doi: 10.1016/j.ypmed.2019.04.027
- [21] Hutchinson, D., Moore, E.A., Breen, C., Burns, L., & Mattick, R.P. (2013). Alcohol use in pregnancy: Prevalence and predictors in the longitudinal study of Australian children. *Drug and alcohol rev.*, 32(5): 475-482. doi: 10.1111/dar.12027
- [22] Scheffers-van Schayck, T., Den Hollander, W., Van Belzen, E., Monshouwer, K., & Tuithof, M. (2019). Monitor Middelengebruik en Zwangerschap 2018. Utrecht: Trimbos-instituut.
- [23] Tuithof, M., Siau, R., Van Dorsseleer, S., & Monshouwer, K. (2017). Factsheet Monitor Zwangerschap en Middelengebruik: Het middelengebruik van moeders en hun partner voor, tijdens en na de zwangerschap. Utrecht: Trimbos-instituut.
- [24] Gezondheidsraad (2005). Risico's van alcoholgebruik bij conceptie, zwangerschap en borstvoeding. Den Haag: Gezondheidsraad.
- [25] Bommele, J., Springvloet, L., Abouri, N., Djoyadhiningrat-Hol, K., Van Laar, M., & Blankers, M. (2020). Stepped-wedge cluster randomised trial of a smoking cessation counselling training programme for midwives treating women with functional health illiteracy and low socioeconomic status (PROMISE): A study protocol. *Trials*, 21(1): 619. doi: 10.1186/s13063-020-04555-0
- [26] Faun, H., van der Velden, M., van der Kemp, S., & Driessen, T. (2017). Alcohol en de kinderschap. Over alcoholgebruik tijdens de zwangerschap. Zoetermeer: Panteia.
- [27] Schouten, F., Tuithof, M., van Hasselt, N., & Voogt, C. (2021). Kennismonitor Alcohol 2020: Onderzoek onder volwassenen naar kennis over alcoholische. Utrecht: Trimbos-instituut.
- [28] Ministerie van Volksgezondheid Welzijn en Sport (2018). Nationaal Preventieakkoord: Naar een gezonder Nederland.
- [29] Stead, L.F., Buitrago, D., Preciado, N., Sanchez, G., Hartmann-Boyce, J., & Lancaster, T. (2013). Physician advice for smoking cessation. *Cochrane Database Syst Rev*; 5. doi: 10.1002/14651858.CD000165.pub4
- [30] Shankaran, S., Lester, B.M., Das, A., Bauer, C.R., Bada, H.S., Lagasse, L., et al. (2007). Impact of maternal substance use during pregnancy on childhood outcome. *Semin Fetal Neonatal Med.*, 12(2):143. doi: 10.1016/j.siny.2007.01.002