

Referenties

- [1] Drake, R.E., Mueser, K.T., & Brunette, M.F. Management of persons with co-occurring severe mental illness and substance use disorder: program implications. *World Psychiatry*. 2007; 6(3): 131-136.
- [2] Haverfield, M.C., Ilgen, M., Schmidt, E., Shelley, A., & Timko, C. Social support networks and symptom severity among patients with co-occurring mental health and substance use disorders. *Community Mental Health*. 2019; 55(5): 768-776. doi:10.1007/s10597-019-00396-7
- [3] Milani, R.M., Nahar, K., Ware, D., Butler, A., Roush, S., Smith, D., Perrino, L., & O'Donnell, J. A qualitative longitudinal study of the first Dual Diagnosis Anonymous (DDA) in the UK. *Advances in dual diagnosis*. 2020; 13(4): 151-167. doi:10.1108/ADD-01-2020-0001
- [4] Chambers, R.A. Adult hippocampal neurogenesis in the pathogenesis of addiction and dual diagnosis disorders. *Drug and alcohol dependence*. 2013; 130(1-3): 1-12. doi:10.1016/j.drugalcdep.2012.12.005
- [5] Drake, R.E., & Brunette, M.F. Complications of severe mental illness related to alcohol and drug use disorders. *Recent developments in alcoholism*. 1998; 14(1): 285-299.
- [6] Mangrum, L.F., Spence, R.T., & Lopez, M. Integrated versus parallel treatment of co-occurring psychiatric and substance use disorders. *Journal of Substance Abuse Treatment*. 2006; 30(1): 79-84. doi:10.1016/j.jsat.2005.10.004
- [7] Neven, A., Kool, N., Bonebakker, A., & Mulder, C. Geïntegreerd behandelen van patienten met dubbele diagnose met integrated dual diagnosis treatment (IDDT): systematische literatuurstudie. *Tijdschrift voor Psychiatrie*. 2018; 60(4): 231-240.
- [8] Kroon, H., & Boyle, P. Integrated Dual Disorder Treatment: Comparing Facilitators and Challenges of Implementation for Ohio and the Netherlands. *International Journal of Mental Health*. 2006; 35(2): 70-88.
- [9] Anthony, W.A. (1993). Recovery from mental illness: The guiding vision of the mental health service system in the 1990s. *Psychosocial Rehabilitation Journal*. 1993; 16(4), 11-23.
- [10] Generieke module Herstelondersteuning. AKWA GGZ, 2017.
- [11] Magura, S., Rosenblum, A., Villano, C.L., Vogel, H.S., Fong, C., & Betzler, T. Dual-focus mutual aid for co-occurring disorders: a quasi-experimental outcome evaluation study. *American Journal of Drug and Alcohol Abuse*. 2008; 34(1): 61-74. doi:10.1080/00952990701764623
- [12] Vogel, H.S., Knight, E., Laudet, A.B., & Magura, S. Double Trouble in Recovery: self-help for people with dual diagnoses. *Psychiatric Rehabilitation Journal*. 1998; 21(4): 356-364.
- [13] Green, C.A., Yarborough, M.T., Polen, M.R., Janoff, S.L., & Yarborough, B.J. Dual Recovery among people with serious mental illnesses and substance problems: a qualitative analysis. *Journal of Dual Diagnosis*. 2015; 11(1): 33-41. doi:10.1080/15504263.2014.975004
- [14] Bogenschutz, M.P., Geppert, C.M., & George, J. The role of twelve-step approaches in dual diagnosis treatment and recovery. *American Journal on Addictions*. 2006; 15(1): 50-60. doi:10.1080/10550490500419060
- [15] Chilton, J., Crone, D.M., & Tyson, P.J. A qualitative study of an integrated group treatment for dual diagnosis service users within a community mental health setting. *International Journal of Mental Health*. 2019; 29(3): 406-413. doi:10.1111/inm.12675
- [16] Ouimette, P., Humphreys, K., Moos, R.H., Finney, J.W., Cronkite, R., & Federman, B. Self-help group participation among substance use disorder patients with posttraumatic stress disorder. *Journal of Substance Abuse Treatment*. 2001; 20(1): 25-32.
- [17] Bogenschutz, M.P. 12-step approaches for the dually diagnosed: mechanisms of change. *Alcoholism: Clinical and Experimental Research*. 2007; 31(30): 64-66. doi:10.1111/j.1530-0277.2007.00496.x
- [18] Chi, F.W., Sterling, S., Campbell, C.I., & Weisner, C. 12-step participation and outcomes over 7 years among adolescent substance use patients with and without psychiatric comorbidity. *Substance abuse*. 2013; 34(1): 33-42. doi:10.1080/08897077.2012.691780
- [19] Herman, S.E., Frank, K.A., Mowbray, C.T., Ribisl, K.M., & Davidson, W.S. Longitudinal effects of integrated treatment on alcohol use for persons with serious mental illness and substance

- use disorder. *Journal of Behavioral Health Services & Research*. 2000; 27(3): 286-302. doi:10.1007/BF02291740
- [20] Kelly, J.F., McKellar, J.D., & Moos, R. Major depression in patients with substance use disorders: relationship to 12-step self-help involvement and substance use outcomes. *Addiction*. 2003; 98(4): 499-508. doi:10.1046/j.1360-0443.2003.t01-1-00294.x
- [21] Tonigan, J.S., Pearson, M.R., Magill, M., & Hagler, K.J. AA attendance and abstinence for dually diagnosed patients: a meta-analytic review. *Addiction*. 2018; 113(11): 1970-1981. doi:10.1111/add.14268
- [22] Timko, C., Sutkowi, A., & Moos, R. Patients with dual diagnoses or substance use disorders only: 12-step group participation and 1-year outcomes. *Substance Use & Misuse*. 2010; 45(4): 613-627. doi:10.3109/10826080903452421
- [23] DiNitto, D.M., Webb, D.K., Rubin, A., Debra, M.-O., & Wambach, K.G. Self-help group meeting attendance among clients with dual diagnosis. *Journal of Psychoactive Drugs*. 2001; 33(3): 263-272. doi: 10.1080/02791072.2001.10400573
- [24] Muusse, C., & van Rooijen, S. De echo van het eigen verhaal: over zelfhulpgroepen bij verslaving en de samenwerking met de ggz en verslavingszorg. 2009. Utrecht: Trimbos-instituut.
- [25] Laudet, A.B., Magura, S., Vogel, H.S., & Knight, E. Support, mutual aid and recovery from dual diagnosis. *Mental Health Journal*. 2000; 36(5): 457-476.
- [26] Noordsy, D.L., Schwab, B., Fox, L., & Drake, R.E. The role of self-help programs in the rehabilitation of persons with severe mental illness and substance use disorders. *Community Mental Health*. 1996; 32(1): 71-81.
- [27] Timko, C., Cronkite, R.C., McKellar, J., Zemore, S., & Moos, R.H. Dually diagnosed patients' benefits of mutual-help groups and the role of social anxiety. *Journal of Substance Abuse Treatment*. 2013; 44(2): 216-223. doi:10.1016/j.jsat.2012.05.007
- [28] Magura, S. Effectiveness of dual focus mutual aid for co-occurring substance use and mental health disorders: a review and synthesis of the "double trouble" in recovery evaluation. *Substance Use & Misuse*. 2009; 43(12-13): 1904-1926. doi:10.1080/10826080802297005
- [29] Roush, S., Monica, C., Carpenter-Song, E., & Drake, R. First-person perspectives on Dual Diagnosis Anonymous (DDA): a qualitative study. *Journal of Dual Diagnosis*. 2015; 11(2): 136-141. doi:10.1080/15504263.2015.1025215
- [30] Hamilton, T., & Samples, P. Twaalf stappen en dubbele diagnose: handboek voor 12-stappen zelfhulpgroepen voor mensen met een verslaving & een psychiatrische aandoening. 2010. Utrecht: Landelijk Expertisecentrum Dubbele Diagnose (LEDD).
- [31] Magura, S., Knight, E.L., Vogel, H.S., Mahmood, D., & Laudet, A.B. Mediators of effectiveness in dual-focus self-help groups. *American Journal of Drug and Alcohol Abuse*. 2003; 29(2): 301-322. doi:10.1081/ADA-120020514
- [32] Magura, S., Laudet, A.B., Mahmood, D., Rosenblum, A., & Knight, E. Adherence to medication regimens and participation in dual-focus self-help groups. *Psychiatric Services*. 2002; 53(3): 310-316. doi:10.1176/appi.ps.53.3.310
- [33] Rosenblum, A., Matusow, H., Fong, C., Vogel, H., Uttaro, T., Moore, T.L., & Magura, S. Efficacy of dual focus mutual aid for persons with mental illness and substance misuse. *Drug and Alcohol Dependence*. 2013; 135(1): 78-87. doi:10.1016/j.drugalcdep.2013.11.012
- [34] Timko, C., Sutkowi, A., Cronkite, R.C., Makin-Byrd, K., & Moos, R.H. Intensive referral to 12-step dual-focused mutual-help groups. *Drug and Alcohol Dependence*. 2011; 118(2-3): 194-201. doi:10.1016/j.drugalcdep.2011.03.019
- [35] Laudet, A.B., Magura, S., Cleland, C.M., Vogel, H.S., Knight, E.L., & Rosenblum, A. The effect of 12-step based fellowship participation on abstinence among dually diagnosed persons: a two-year longitudinal study. *Journal of Psychoactive Drugs*. 2004; 36(2): 207-216. doi:10.1080/02791072.2004.10399731
- [36] Magura, S., Laudet, A.B., Mahmood, D., Rosenblum, A., Vogel, H.S., & Knight, E.L. Role of self-help processes in achieving abstinence among dually diagnosed persons. *Addictive Behaviors*. 2003; 28(3): 399-413. doi:10.1016/S0306-4603(01)00278-7

- [37] Pallaveshi, L., Balachandra, K., Subramanian, P., & Runick, A. Peer-led and professional-led group interventions for people with co-occurring disorders: a qualitative study. *Mental Health Journal*. 2014; 50(4): 388-394. doi:10.1007/s10597-013-9612-8
- [38] Bogenschutz, M.P., Rice, S.L., Tonigan, J.S., Vogel, H.S., Nowinski, J., Hume, D., & Arenella, P.B. 12-step facilitation for the dually diagnosed: a randomized clinical trial. *Journal of Substance Abuse Treatment*. 2014; 46(4): 403-411. doi:10.1016/j.jsat.2013.12.009