

## Middelengebruik in de verschillende fasen van psychose

Sanne van der Heijden & Jentien Vermeulen

- [1] Volkow ND. Substance use disorders in schizophrenia--clinical implications of comorbidity. *Schizophr Bull.* 2009;35(3):469-72.
- [2] WHO. WHO global report on trends in prevalence of tobacco use 2000-2025; 2020. 2022.
- [3] Arseneault L, Cannon M, Witton J, Murray RM. Causal association between cannabis and psychosis: examination of the evidence. *Brit J Psychiat.* 2004;184(2):110-7.
- [4] Hartz SM, Pato CN, Medeiros H, Cavazos-Rehg P, Sobell JL, Knowles JA, et al. Comorbidity of Severe Psychotic Disorders With Measures of Substance Use. *JAMA Psychiatry.* 2014;71(3):248-54.
- [5] Lahteenvuo M, Batalla A, Luykx JJ, Mittendorfer-Rutz E, Tanskanen A, Tiihonen J, Taipale H. Morbidity and mortality in schizophrenia with comorbid substance use disorders. *Acta Psychiatr Scand.* 2021.
- [6] Alvarez-Jimenez M, Priede A, Hetrick SE, Bendall S, Killackey E, Parker AG, et al. Risk factors for relapse following treatment for first episode psychosis: A systematic review and meta-analysis of longitudinal studies. *Schizophrenia Research.* 2012;139(1):116-28.
- [7] Carney R, Yung A, Amminger G, Bradshaw T, Glozier N, Hermens D, et al. Substance use in youth at risk for psychosis. *Schizophrenia research.* 2016;181.
- [8] Ward HB, Lawson MT, Addington J, Bearden CE, Cadenhead KS, Cannon TD, et al. Tobacco use and psychosis risk in persons at clinical high risk. *Early Interv Psychiatry.* 2019;13(5):1173-81.
- [9] Schizophrenia ENoNNsG-Eli, van Os J, Rutten BP, Myin-Germeys I, Delespaul P, Viechtbauer W, et al. Identifying gene-environment interactions in schizophrenia: contemporary challenges for integrated, large-scale investigations. *Schizophr Bull.* 2014;40(4):729-36.
- [10] Korver N, Quee PJ, Boos HB, Simons CJ, de Haan L. Genetic Risk and Outcome of Psychosis (GROUP), a multi-site longitudinal cohort study focused on gene-environment interaction: objectives, sample characteristics, recruitment and assessment methods. *Int J Methods Psychiatr Res.* 2012;21(3):205-21.
- [11] Segeren M, Lauriks S, Kikkert M, Heering J, Lommerse N, van Husen G, Verhoeff A. Deinstitutionalization from the perspective of community-dwelling adults with a severe mental illness in Amsterdam: a cohort study protocol. *BMC Public Health.* 2022;22(1):950.
- [12] Dariotis J, Chen F. Stress Coping Strategies as Mediators: Toward a Better Understanding of Sexual, Substance, and Delinquent Behavior-Related Risk-Taking among Transition-Aged Youth. *Deviant Behavior.* 2020:1-18.
- [13] Moylan S, Jacka FN, Pasco JA, Berk M. Cigarette smoking, nicotine dependence and anxiety disorders: a systematic review of population-based, epidemiological studies. *BMC Med.* 2012;10:123.

- [14] Aronson KR, Almeida DM, Stawski RS, Klein LC, Kozlowski LT. Smoking is Associated with Worse Mood on Stressful Days: Results from a National Diary Study. *Annals of Behavioral Medicine*. 2008;36(3):259-69.
- [15] Vermeulen J, Schirmbeck F, Blankers M, Van Tricht M, Bruggeman R, Van den Brink W, De Haan L. Association Between Smoking Behavior and Cognitive Functioning in Patients With Psychosis, Siblings, and Healthy Control Subjects: Results From a Prospective 6-Year Follow-Up Study. *American Journal of Psychiatry*. 2018;0(0):appi.ajp.2018.18010069.
- [16] Vermeulen J, Schirmbeck F, Blankers M, van Tricht M, van den Brink W, de Haan L, et al. Smoking, symptoms, and quality of life in patients with psychosis, siblings, and healthy controls: a prospective, longitudinal cohort study. *The Lancet Psychiatry*. 2019;6(1):25-34.
- [17] Wootton RE, Richmond RC, Stuijzand BG, Lawn RB, Sallis HM, Taylor GMJ, et al. Evidence for causal effects of lifetime smoking on risk for depression and schizophrenia: a Mendelian randomisation study. *Psychol Med*. 2019:1-9.
- [18] Gogos A, Skokou M, Ferentinou E, Gourzis P. Nicotine consumption during the prodromal phase of schizophrenia - a review of the literature. *Neuropsychiatr Dis Treat*. 2019;15:2943-58.
- [19] Moritz S, Ludtke T, Westermann S, Hermeneit J, Watroba J, Lincoln TM. Dysfunctional coping with stress in psychosis. An investigation with the Maladaptive and Adaptive Coping Styles (MAX) questionnaire. *Schizophr Res*. 2016;175(1-3):129-35.
- [20] Thornton LK, Baker AL, Lewin TJ, Kay-Lambkin FJ, Kavanagh D, Richmond R, et al. Reasons for substance use among people with mental disorders. *Addict Behav*. 2012;37(4):427-34.
- [21] Taylor GM, Lindson N, Farley A, Leinberger-Jabari A, Sawyer K, Te Water Naude R, et al. Smoking cessation for improving mental health. *Cochrane Database Syst Rev*. 2021;3:CD013522.
- [22] Provencal N, Binder EB. The neurobiological effects of stress as contributors to psychiatric disorders: focus on epigenetics. *Current Opinion in Neurobiology*. 2015;30:31-7.
- [23] Mazzone P, Tierney W, Hossain M, Puvenna V, Janigro D, Cucullo L. Pathophysiological impact of cigarette smoke exposure on the cerebrovascular system with a focus on the blood-brain barrier: expanding the awareness of smoking toxicity in an underappreciated area. *International journal of environmental research and public health*. 2010;7(12):4111-26.
- [24] Schaan VK, Schulz A, Rubel JA, Bernstein M, Domes G, Schachinger H, Vogele C. Childhood Trauma Affects Stress-Related Interoceptive Accuracy. *Frontiers in Psychiatry*. 2019;10.
- [25] Ferraro L, La Cascia C, Quattrone D, Sideli L, Matranga D, Capuccio V, et al. Premorbid Adjustment and IQ in Patients With First-Episode Psychosis: A Multisite Case-Control Study of Their Relationship With Cannabis Use. *Schizophrenia Bulletin*. 2019;46(3):517-29.
- [26] Salyers MP, Mueser KT. Social functioning, psychopathology, and medication side effects in relation to substance use and abuse in schizophrenia. *Schizophr Res*. 2001;48(1):109-23.
- [27] Gneiting U, Schmitz HP. Comparing global alcohol and tobacco control efforts: network formation and evolution in international health governance. *Health Policy and Planning*. 2016;31(suppl\_1):i98-i109.

[28] McGinty EE, Goldman HH, Pescosolido B, Barry CL. Portraying mental illness and drug addiction as treatable health conditions: Effects of a randomized experiment on stigma and discrimination. *Social Science & Medicine*. 2015;126:73-85.

[29] Taylor G, McNeill A, Girling A, Farley A, Lindson-Hawley N, Aveyard P. Change in mental health after smoking cessation: systematic review and meta-analysis. *BMJ*. 2014;348:g1151.

[30] Zeng LN, Zong QQ, Zhang L, Feng Y, Ng CH, Ungvari GS, et al. Worldwide prevalence of smoking cessation in schizophrenia patients: A meta-analysis of comparative and observational studies. *Asian J Psychiatr*. 2020;54:102190.