

Referenties

Associatie tussen alcoholconsumptie en coping als drinkmotief in een klinische populatie

Matthieu Lingier, Michiel van Kernebeek, Nathalie Vanderbruggen & Cleo L. Crunelle

- [1] Griswold, M.G., Fullman, N., Hawley, C., Arian, N., Zimsen, S.R.M., Tymeson, H.D., e.a. (2018). Alcohol use and burden for 195 countries and territories, 1990-2016: A systematic analysis for the Global Burden of Disease Study 2016. *The Lancet*, 392(10152): 1015–35.
- [2] WHO (2018). Global status report on alcohol and health 2018.
- [3] Magill, M., & Ray, L.A. (2009). Cognitive-behavioral treatment with adult alcohol and illicit drug users: a meta-analysis of randomized controlled trials. *J Stud Alcohol Drugs*, 70(4): 516–27.
- [4] Litten, R.Z., Falk, D.E., Ryan, M.L., & Fertig, J.B. (2016). Discovery, Development, and Adoption of Medications to Treat Alcohol Use Disorder: Goals for the Phases of Medications Development. *Alcohol Clin Exp Res.*, 40(7): 1368–79.
- [5] Bresin, K., & Mekawi, Y. (2021). The “Why” of Drinking Matters: A Meta-Analysis of the Association Between Drinking Motives and Drinking Outcomes. *Alcohol Clin Exp Res.*, 45(1): 38–50.
- [6] Cooper, M.L. (1994). Motivations for alcohol use among adolescents: Development and validation of a four-factor model. *Psychol Assess.*, 6(2): 117–28.
- [7] Cooper, M.L., Kuntsche, E., Levitt, A., Barber, L.L., & Wolf, S. (2015). Motivational Models of Substance Use. Sher, K.J. (ed.). Vol. 1. Oxford University Press.
- [8] Cooper, M.L., Frone, M.R., Russell, M., & Mudar, P. (1995). Drinking to regulate positive and negative emotions: a motivational model of alcohol use. *J Pers Soc Psychol.*, 69(5): 990–1005.
- [9] Mezquita, L., Stewart, S.H., Ibáñez, M.I., Ruipérez, M.A., Villa, H., Moya, J., e.a. (2011). Drinking Motives in Clinical and General Populations. *Eur Addict Res.*, 17(5): 250–61.
- [10] Baumeister, R.F., & Scher, S.J. (1988). Self-defeating behavior patterns among normal individuals: Review and analysis of common self-destructive tendencies. *Psychol Bull.*, 104(1): 3–22.
- [11] Hayes, S.C., Strosahl, K., Wilson, K.G., Bissett, R.T., Pistorello, J., Toarmino, D., e.a. (2004). Measuring experiential avoidance: A preliminary test of a working model. *Psychol Rec.*, 54(4): 553–78.
- [12] Cornelius, J.R., Bukstein, O., Salloum, I., & Clark, D. (2002). Alcohol and Psychiatric Comorbidity. In: *Recent Developments in Alcoholism*. Boston, MA: Springer US, p. 361–74.
- [13] Nehlin, C., & Öster, C. (2019). Measuring drinking motives in undergraduates: An exploration of the Drinking Motives Questionnaire-Revised in Swedish students. *Subst Abuse Treat Prev Policy*, 14(1).
- [14] Kuntsche, E., Knibbe, R., Gmel, G., & Engels, R. (2005). Why do young people drink? A review of drinking motives. *Clin Psychol Rev.*, 25(7): 841–61.

- [15] Blevins, C.E., & Stephens, R.S. (2016). The impact of motives-related feedback on drinking to cope among college students. *Addictive behaviors*, 58: 68–73.
- [16] Anker, J.J., Forbes, M.K., Almquist, Z.W., Menk, J.S., Thuras, P., Unruh, A.S., e.a. (2017). A network approach to modeling comorbid internalizing and alcohol use disorders. *J Abnorm Psychol.*, 126(3): 325–39.
- [17] von Elm, E., Altman, D.G., Egger, M., Pocock, S.J., Gøtzsche, P.C., & Vandebroucke, J.P. (2007). The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: Guidelines for Reporting Observational Studies. *Ann Intern Med.*, 147(8): 573.
- [18] Babor, T.F., Higgins-Biddle, J.C., Saunders, J.B., & Monteiro, M.G. (2001). World Health Organization.
- [19] Saunders, J.B., Aasland, O.G., Babor, T.F., de la Fuente, J.R., & Grant, M. (1993). Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption--II. *Addiction*, 88(6): 791–804.
- [20] Kuntsche, E., & Kuntsche, S. (2009). Development and validation of the drinking motive questionnaire revised short form (DMQ-R SF). *Journal of Clinical Child and Adolescent Psychology*, 38(6): 899–908.
- [21] Kessler, R.C. (2004). The epidemiology of dual diagnosis. *Biol Psychiatry*, 56(10): 730–7.
- [22] Nehlin, C., Grönbladh, L., Fredriksson, A., & Jansson, L. (2013). Alcohol and drug use, smoking, and gambling among psychiatric outpatients: a 1-year prevalence study. *Subst Abus.*, 34(2): 162–8.
- [23] Kushner, M.G., Maurer, E.W., Thuras, P., Donahue, C., Frye, B., Menary, K.R., e.a. (2013). Hybrid cognitive behavioral therapy versus relaxation training for co-occurring anxiety and alcohol disorder: a randomized clinical trial. *J Consult Clin Psychol.*, 81(3): 429–42.
- [24] Carpenter, K.M., & Hasin, D. (1998). A prospective evaluation of the relationship between reasons for drinking and dsm-iv alcohol-use disorders. *Addictive Behaviors*, 23(1): 41–6.
- [25] Di Chiara, G. (1997). Alcohol and dopamine. *Alcohol Health Res World*, 21(2): 108–14.
- [26] Layard, R., & Sachs, J. (2013). World happiness report 2013.
- [27] Pernanen, K. (2001). Consequences of drinking to friends and the close social environment. In: *Mapping the Social Consequences of Alcohol Consumption*. Dordrecht: Springer Netherlands, p. 53–65.
- [28] Van Damme, J., Maes, L., Clays, E., Rosiers, J.F.M.T., Van Hal, G., & Hublet, A. (2013). Social motives for drinking in students should not be neglected in efforts to decrease problematic drinking. *Health Educ Res.*, 28(4): 640–50.
- [29] Monti, P.M., & Rohsenow, D.J. (1999). Coping-skills training and cue-exposure therapy in the treatment of alcoholism. *Alcohol Res Health*, 23(2): 107–15.